

HARD THERMAL LOOPS AND QCD THERMODYNAMICS*

A. REBHAN

Institut für Theoretische Physik
Technische Universität Wien
Wiedner Hauptstr. 8-10, A-1040 Vienna, Austria

Abstract: These lectures give an introduction to thermal perturbation theory, hard thermal loops, and their use in a nonperturbative, approximately self-consistent resummation of the thermodynamical potentials of quantum chromodynamics.

1 Introduction

At sufficiently high temperature and/or density, quantum chromodynamics (QCD) should become accessible by perturbative methods due to asymptotic freedom. Thermal perturbation theory is however a surprisingly intricate subject, and it was only during the last decade that the necessary methodology has been developed, and this development is not yet completely finished. An important milestone was the hard-thermal-loop (HTL) resummation programme introduced in particular by Braaten and Pisarski [23] which generalized the static ring resummations that were known to be required in the calculation of thermo-*static* quantities since long [40, 1] to dynamic quantities such as quasi-particle properties as well as production and loss rates.

There are, however, well-known limitations of a fundamental nature which present an impenetrable barrier to perturbation theory at a certain order of the coupling (depending on the quantity under consideration), caused by the inherently nonperturbative chromo-magnetostatic sector of nonabelian gauge theories [68, 61, 42]. This does not mean that thermal perturbation theory is completely futile, though, but rather that at particular points certain nonperturbative input is required in addition [20], for example from lattice calculations of the inherently nonperturbative 3-d Yang-Mills theory describing the self-interactions of chromo-magnetostatic fields. There are in fact even more (less well-known) limitations from collinear singularities that occur in real-time quantities involving external light-like momenta such as the production rate of real photons from a QCD plasma [11, 8], which require improved resummations [32] and/or nonperturbative input.

*Lectures given at the NATO Advanced Study Institute “QCD perspectives on hot and dense matter”, August 6–18, 2001, in Cargèse, Corsica, France

But even in those cases where thermal perturbation theory has not yet run into one of those barriers, such as the landmark three-loop calculation of the free energy of QCD to order $\alpha_s^{5/2}$ by Arnold, Zhai, and others [7, 21], there are severe problems caused by extremely poor convergence and strong renormalization scheme dependences which seem to render quantitative predictions possible only beyond preposterously high temperatures. While some improvement seemed possible through tricks like Padé approximations [47, 43], the sad conclusion appeared to be that thermal perturbation theory was not applicable in QCD at temperatures of practical interest [22].

Recently, however, it became clear that the problem of poor convergence is not specific to QCD, but arises already in such simple theories as massless scalar φ^4 theory [30] and that alternative resummations can be found that greatly improve the apparent convergence [46, 4]. In these lectures, after an introduction to hard thermal loops, I will present an approach to the problem of calculating thermodynamical quantities in QCD in an *approximately self-consistent* resummation of hard thermal loops and their next-to-leading order corrections [15, 16, 17, 18] that appears to work well down to temperatures a few times the deconfinement transition temperature and that suggests that at such temperatures the still strongly interacting QCD may (at least in certain cases) be adequately described by weakly interacting HTL quasiparticles after all.

2 Thermal field theory

The Feynman rules of thermal field theory [45, 57] are most easily formulated in the imaginary-time (ITF) or Matsubara formalism. The statistical density operator $e^{-\beta H}$ is then equivalent to a time evolution operator over an imaginary time interval of length β , the inverse temperature T^{-1} , and the traces in $\langle A \rangle = \text{Tr}[e^{-\beta H} A]$ require (anti-)periodic boundary conditions at the ends of the imaginary time interval for bosons (fermions). This gives rise to discrete imaginary (Matsubara) frequencies when going to momentum space, so that the only change in the Feynman rules is in the replacement

$$\int \frac{d^4 k}{i(2\pi)^4} \rightarrow \beta^{-1} \sum_{\nu} \int \frac{d^3 k}{(2\pi)^3}, \quad i(2\pi)^4 \delta^4(k) \rightarrow \beta(2\pi)^3 \delta_{\nu,0} \delta^3(k). \quad (1)$$

with

$$k_0 \rightarrow 2\pi i T \nu, \quad \nu \in \begin{cases} \mathbb{Z} & \text{bosons} \\ \mathbb{Z} - \frac{1}{2} & \text{fermions} \end{cases}. \quad (2)$$

Since it is usually hard to evaluate the resulting sums directly, they are best turned into integrals again by writing, in the bosonic case,

$$T \sum_{n=-\infty}^{\infty} f(k_0 = 2\pi i n T) = \frac{T}{2\pi i} \oint_{\mathcal{C}} dk_0 f(k_0) \frac{\beta}{2} \coth \frac{\beta k_0}{2}$$

$$= \underbrace{\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} f(k_0)}_{\text{vacuum contribution}} + \underbrace{\frac{1}{2\pi i} \int_{-i\infty+\epsilon}^{i\infty+\epsilon} [f(k_0) + f(-k_0)] \frac{1}{e^{\beta k_0} - 1}}_{\text{thermal contribution}} \quad (3)$$

for a function $f(k_0)$ that is regular for $k_0 \in i\mathbb{R}$ and where $\mathcal{C}$ is a contour encircling only the poles of $\coth$ at the location of the Matsubara frequencies. (In the fermionic case, an analogous formula can be easily obtained by using $\tanh$ in place of $\coth$, which in the right-hand side leads to minus the Fermi-Dirac distribution in place of the Bose-Einstein one.)

This evaluation through contour integrals leads to a nice separation into a (Wick rotated) $T = 0$ contribution, and a purely thermal one, which vanishes for $\beta \rightarrow \infty$, i.e. $T \rightarrow 0$. There exists also a formulation directly in real time (Schwinger-Keldysh and variants thereof) [56] where this separation is already conspicuous in the Feynman rules. However, this requires a doubling of fields, a 2×2 matrix structure for propagators and even more components for n -point vertex functions, which in fact correspond to the many possibilities of analytic continuation to real frequencies (in particular if there are several independent external frequencies) [49].

A simple case that leads to a thermal contribution in (3) is the frequently occurring one that there is a simple pole in $f(k_0)$ at $k_0 = \pm E$ with $E > 0$. Then one has

$$T \sum_n \frac{1}{k_0 \pm E} = \pm (-1)^\sigma n_\sigma(E) + \text{vacuum contributions}, \quad (4)$$

where $n_\sigma(E) = [e^{\beta E} - \sigma]^{-1}$ and $\sigma = +1$ for bosons, -1 for fermions.

3 Hard thermal loops

The simplest example of a hard thermal loop is given by the one-loop self-energy diagram in a scalar $g\varphi^4$ theory. This is a tadpole diagram, independent of external momenta:

$$\Pi = \frac{4!g^2}{2} \sum_K \frac{-1}{K^2 - m^2} \quad (5)$$

where we have introduced the notation $\sum_K = T \sum_n \int \frac{d^3 k}{(2\pi)^3}$, $K^\mu = (k^0, k^m)$.

With $E = \sqrt{k^2 + m^2}$, its thermal contribution is easily evaluated as

$$\begin{aligned} \sum_K \frac{-1}{K^2 - m^2} &= \sum_K \frac{1}{2E} \left(\frac{1}{k_0 - E} - \frac{1}{k_0 + E} \right) \\ &= \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2E} 2n(E) = \int_0^\infty \frac{dk k^2}{2\pi^2 E} n(E). \end{aligned} \quad (6)$$

(The vacuum contribution is removed by standard ($T=0$) renormalization.) Without the Bose-Einstein factor, this integral would be quadratically divergent. Thanks to the former, it is finite, but dominated by momenta $k \sim T$. If

$T \gg m$, the mass terms of the $T=0$ theory can be neglected, and the leading contribution to the self-energy is given by a self-energy contribution proportional to T^2 , which is called a *hard thermal loop* (HTL):

$$\hat{\Pi} = \frac{4!g^2}{2} \sum_K \frac{-1}{K^2} = g^2 \frac{6}{\pi^2} \int_0^\infty dk k n(k) = g^2 T^2, \quad (7)$$

where the hat is a reminder of the HTL approximation – in this case the neglect of the bare mass m .

$\hat{\Pi}$ clearly corresponds to a mass term for scalar excitations generated by interactions with the particles in the heat bath. It should be understood, however, that this *thermal mass* is qualitatively different from ordinary masses. In particular, one can show that unlike ordinary masses it does not spoil conformal invariance (if any) as it does not contribute to the trace of the energy momentum tensor [63].

This example for a HTL is, however, deceptively simple. In general, thermal masses are not constant but depend on momentum, that is, they correspond to nonlocal terms in an effective action.

The thermal masses generated for gauge fields are of this form. Indeed, a constant mass term would violate gauge invariance. Let us consider as a simple gauge theory example the case of scalar electrodynamics with covariant gauge fixing,

$$\mathcal{L} = (D_\mu \phi)^* D^\mu \phi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2\alpha} (\partial_\mu A^\mu)^2 \quad (8)$$

where $D_\mu = \partial_\mu + ieA_\mu$. The tree-level propagator reads $G_{\mu\nu}^0 = g_{\mu\nu}/K^2 + (\alpha - 1)K_\mu K_\nu/K^4$ and this will receive thermal corrections through the photon polarization tensor, which in momentum space reads

$$\Pi_{\mu\nu}(K) = e^2 \sum_P \left[\frac{(2P - K)_\mu (2P - K)_\nu}{P^2 (P - K)^2} - \frac{2g_{\mu\nu}}{P^2} \right]. \quad (9)$$

The last term is from the seagull diagram, which is essentially the same as the tadpole diagram before. The first term is however dependent on the external momentum, and there is no reason to expect a Lorentz invariant form, because the heat bath is singling out a preferred frame of reference.

There are in fact four symmetric tensors that can be built from the available quantities $g_{\mu\nu}$, K_μ , and the four-velocity of the heat bath which we have tacitly chosen as $U_\mu = \delta_\mu^0$.

In electrodynamics, the polarization tensor has to be transverse, $K^\mu \Pi_{\mu\nu} \equiv 0$. This additional requirement still leaves two possible tensors, for from U and K one can build the transverse vector $\tilde{U}_\mu = K^2 U_\mu - (U \cdot K) K_\mu$. We choose them as

$$A_{\mu\nu} = g_{\mu\nu} - K_\mu K_\nu / K^2 - B_{\mu\nu}, \quad B_{\mu\nu} = \tilde{U}_\mu \tilde{U}_\nu / \tilde{U}^2. \quad (10)$$

$A_{\mu\nu}$ can easily be shown to have vanishing components $A_{0\nu} = 0 = A_{\mu 0}$, whereas $A_{ij} = -\delta_{ij} + k_i k_j / k^2$, so this is a projection onto spatially transverse momenta; $B_{\mu\nu}$ is a projector orthogonal to $A_{\mu\nu}$.

In nonabelian gauge theories like QCD, one generally has a more complicated structure. Then one needs two more, nontransverse, tensors, which one may choose as

$$C_{\mu\nu} = \frac{\tilde{U}_\mu K_\nu + K_\mu \tilde{U}_\nu}{\sqrt{2}K^2 k}, \quad D_{\mu\nu} = \frac{K_\mu K_\nu}{K^2}. \quad (11)$$

Decomposing $\Pi_{\mu\nu} = -\Pi_t A_{\mu\nu} - \Pi_\ell B_{\mu\nu} - \Pi_c C_{\mu\nu} - \Pi_d D_{\mu\nu}$ and

$$G^{\mu\nu} \equiv [G_0^{-1} + \Pi]^{-1\mu\nu} = \Delta_t A_{\mu\nu} + \Delta_\ell B_{\mu\nu} + \Delta_c C_{\mu\nu} + \Delta_d D_{\mu\nu} \quad (12)$$

one has

$$\Delta_t = [K^2 - \Pi_t]^{-1}, \quad \Delta_\ell = [K^2 - \Pi_\ell + \alpha \frac{\Pi_c^2}{K^2 - \alpha \Pi_d}]^{-1}, \quad (13)$$

$$\Delta_c = \alpha \Pi_c [K^2 - \alpha \Pi_d]^{-1}, \quad \Delta_d = \frac{\alpha(K^2 - \Pi_\ell)}{K^2 - \alpha \Pi_d} \Delta_\ell. \quad (14)$$

In view of the explicit gauge parameter dependences (there are in fact more hidden within the structure functions of Π), it is remarkable that one can prove that the singularities of Δ_t and Δ_ℓ are gauge-fixing independent [51, 52, 74].

In electrodynamics this situation is much simpler (unless one introduces nonlinear gauge fixing): $\Pi_{\mu\nu}$ is both transverse and completely gauge parameter independent.

Because of transversality, which is easily verified for (9), there are only two independent components, e.g. Π_μ^μ and Π_{00} . The former reads

$$\Pi_\mu^\mu(K) = e^2 \sum_P \left[\frac{-4}{P^2} - \frac{K^2}{P^2(P-K)^2} \right] = 4e^2 \sum_P \frac{-1}{P^2} + O(T) = \frac{e^2 T^2}{3} + O(T) \quad (15)$$

where the term proportional to K^2 does not constitute a hard thermal loop, because its integrand does not involve a quadratic divergence in its vacuum piece. For $k_0, k \ll T$, $\Pi_\mu^\mu(K) \sim \hat{\Pi}_\mu^\mu = e^2 T^2/3$. Notice, however, that this ‘HTL approximation’ remains valid even for $k_0, k \sim T$ as long as $K^2 \ll T^2$.

The other component, Π_{00} is more complicated to evaluate. Its HTL piece, which is contained in

$$\Pi_{00}(K) = e^2 \sum_P \left[\frac{4p_0(p_0 - k_0)}{P^2(P-K)^2} - \frac{2}{P^2} \right], \quad (16)$$

can however be extracted rather easily using

$$\frac{p_0}{P^2} = \frac{1}{2} \left(\frac{1}{p_0 - p} + \frac{1}{p_0 + p} \right), \quad \frac{1}{p_0 + X} \frac{1}{p_0 + Y} = \frac{1}{X - Y} \left(\frac{1}{p_0 + Y} - \frac{1}{p_0 + X} \right)$$

which gives a number of terms of the form $\frac{1}{X-Y}(n(Y) - n(X))$ after summing over the Matsubara frequencies. Now, HTL contributions arise from $p \sim T$, and

for $k_0, k \ll T$ one can approximate the energies X and Y by $\pm p$, except when two hard energies form a soft difference. This gives

$$\Pi_{00}(K) \sim e^2 \int \frac{d^3p}{(2\pi)^3} \left\{ \frac{1}{k_0 + p - |p - k|} [n(|p - k|) - n(p)] + \frac{1}{k_0 - p + |p - k|} [-n(|p - k|) + n(p)] \right\} \quad (17)$$

where the tadpole-like last term in (16) has cancelled against terms where the energy denominators contain the sum of two hard energies.

Since $k \ll p \sim T$, we have $n(|p - k|) - n(p) \sim n'(p)[|p - k| - p]$ and $|p - k| - p \sim \vec{p} \cdot \vec{k}/p \equiv zk$, so the HTL piece of Π_{00} is finally given by

$$\begin{aligned} \hat{\Pi}_{00}(K) &= 2e^2 \int \frac{p^2 dp}{(2\pi)^2} n'(p) \int_{-1}^1 dz \left[-1 + \frac{k_0}{k_0 - zk} \right] \\ &= \frac{e^2 T^2}{3} \left[1 - \frac{k_0}{2k} \ln \frac{k_0 + k}{k_0 - k} \right]. \end{aligned} \quad (18)$$

Originally, k_0 was restricted to Matsubara frequencies. In order to allow for soft $k_0 \ll T$ without being restricted to the zero mode, we in fact need analytic continuation, e.g. $k_0 \rightarrow \omega + i\epsilon$ for retarded boundary conditions, and this defines the cut of the logarithm in (18).

The above results for $\hat{\Pi}_\mu^\mu$ and $\hat{\Pi}_{00}(K)$ are actually universal. They have the same form in nonabelian gauge theories, only the overall coefficient differs. In $SU(N)$ gauge theories with N_f quark flavors one just needs to replace [44, 79] $e^2 \rightarrow g^2(N + N_f/2)$. They also retain their form in the presence of a nonvanishing chemical potential μ_f , which leads to the replacement of $T^2 \rightarrow T^2 + 3\mu_f^2/\pi^2$ in the contributions $\propto N_f$. In the HTL approximation, there are moreover no nontransverse contributions¹ so the HTL gauge boson propagator involves two independent branches determined by $\hat{\Pi}_\ell = -\hat{\Pi}_{00}K^2/k^2$ and $\hat{\Pi}_t = (\hat{\Pi}_\mu^\mu - \hat{\Pi}_\ell)/2$.

The poles of the propagators Δ_t and Δ_ℓ determine the dispersion laws of two sorts of quasiparticles, which in contrast to the scalar φ^4 example are not given by simple mass hyperboloids. These are displayed in Fig. 1 in a plot of $\omega_{t,\ell}(k)$ in quadratic scales (where a relativistic mass hyperboloid would show up as a straight line parallel to the light-cone).

Above a common plasma frequency $\omega_{\text{pl.}} = eT/3$, there are propagating modes, which for large momenta in the transverse branch tend to a mass hyperboloid with asymptotic mass $m_\infty^2 = \frac{3}{2}\omega_{\text{pl.}}^2$, and in branch ℓ approach the light-cone exponentially with exponentially vanishing residue. Indeed, this mode does not have an analogue in the $T=0$ theory but is a purely collective phenomenon, so it has to disappear from the spectrum as $k \rightarrow \infty$. The spatially transverse mode, on the other hand, represents quasiparticles that are in-medium versions of the physical polarisations of gauge bosons.

¹At order kT one has however $\Pi_c \neq 0$ for $\alpha \neq 1$ in the nonabelian case, and $\Pi_d \neq 0$ at two-loop order whenever $\Pi_c \neq 0$.




Figure 1: The location of the zeros of $\Delta_t^{\text{HTL}-1}$ (spatially transverse gauge bosons) and of $\Delta_\ell^{\text{HTL}-1}$ (longitudinal plasmons) in quadratic scales such as to show propagating modes and screening phenomena on one plot.

For $\omega < \omega_{\text{pl}}$, $|\mathbf{k}|$ is the inverse screening length, which in the static limit vanishes for mode t (absence of magnetostatic screening), but reaches the Debye mass, $\hat{m}_D^2 = 3\omega_{\text{pl}}^2$, for mode ℓ (electrostatic screening). A vanishing magnetic screening mass is required by gauge invariance in abelian gauge theories [34, 14], but not in the nonabelian case. In fact, lattice simulations of gauge fixed propagators in nonabelian theories do find a screening behaviour in the transverse sector, however the corresponding singularity is certainly quite different from a simple pole [28].

For $\omega^2 < k^2$, there is a large imaginary part $\sim e^2 T^2$ from (18) which prevents the appearance of poles in this region. This imaginary part corresponds to the possibility of Landau damping, which is the transfer of energy from soft fields to hard plasma constituents moving in phase with the field [60, 13] and is an important part of the spectral density of HTL propagators. At higher, subleading orders of perturbation theory, it is, however, not protected against gauge dependences in nonabelian gauge theories.

To complete the discussion of HTL's in scalar electrodynamics, let us also consider briefly the scalar self-energy. In contrast to φ^4 theory, this is now a nonlocal quantity. Nevertheless, the HTL part is still a constant thermal mass as given by the first term in

$$\Xi = \frac{e^2 T^2}{4} + e^2 K^2 \sum_P \left[\frac{3 - \alpha}{P^2(P - K)^2} + \frac{2(\alpha - 1)K \cdot P}{P^4(P - K)^2} \right]. \quad (19)$$

The other terms are not proportional to T^2 because the integrands do not grow sufficiently at large momenta. They are even gauge parameter dependent, in

contrast to the HTL piece. Notice that these gauge dependent terms vanish on the lowest-order mass shell $K^2 = 0$.

In QCD (and already in spinor QED), we also need to consider the fermion self-energies [48, 80]. In the ultrarelativistic high-temperature limit, bare masses can be neglected. In this case the fermion self-energy can be parametrized by

$$\Sigma(K) = a(K)\gamma^0 + b(K)\vec{k} \cdot \vec{\gamma}/k. \quad (20)$$

Using the same methods as above one easily computes the HTL contributions in

$$\frac{1}{4}\text{tr } \not{K}\Sigma = \omega a + kb = \frac{e^2 T^2}{8} \equiv \hat{M}^2, \quad (21)$$

$$\frac{1}{4}\text{tr } \gamma^0 \Sigma = a = \frac{e^2 T^2}{16k} \ln \frac{\omega + k}{\omega - k}, \quad (22)$$

where in nonabelian gauge theories now $e^2 \rightarrow g^2(N^2 - 1)/(2N)$.

The structure of the HTL fermion propagator is

$$S(K) = \frac{1}{2} \left(\gamma^0 + \frac{\vec{k} \cdot \vec{\gamma}}{k} \right) \Delta_+ + \frac{1}{2} \left(\gamma^0 - \frac{\vec{k} \cdot \vec{\gamma}}{k} \right) \Delta_- \quad (23)$$

with $\Delta_{\pm}^{-1} = -[\omega \mp (k + \Sigma_{\pm})]$ and $\Sigma_{\pm} \equiv b \pm a$. The two branches correspond to spinors whose chirality is equal (+) or opposite (−) to their helicity.

The additional collective modes of branch (−) (“plasminos”) have a curious minimum of ω at $\omega/\hat{M} \approx 0.93$ and $|\vec{k}|/\hat{M} \approx 0.41$ and approach the light-cone for large momenta, but with exponentially vanishing residue. The regular branch approaches a mass hyperboloid (in Fig. 2 a straight line parallel to the diagonal) with asymptotic mass $\sqrt{2}\hat{M}$. Again, for space-like momenta, $K^2 < 0$, there is a large imaginary part corresponding to Landau damping, which now corresponds to the transmutation of soft fermionic fields together with hard fermionic (bosonic) plasma constituents into hard bosonic (fermionic) ones.

4 Standard HTL perturbation theory

When writing down dressed propagators, as we have done above in the discussion of the spectrum of thermal quasiparticles, we have already performed a resummation of infinitely many loops—the geometric series of self-energy insertions according to Dyson’s equation, which in the scalar φ^4 theory is simply

$$\frac{-1}{K^2 - (gT)^2} = \frac{-1}{K^2} \left(1 + g^2 \frac{T^2}{K^2} + g^4 \frac{T^4}{K^4} + \dots \right). \quad (24)$$

Clearly, the perturbative version is useful only when $K^2 \gg (gT)^2$. In particular, it fails for $k_0, k \sim gT$, the scale where collective phenomena transform the familiar quanta into quasiparticles.




Figure 2: The location of the zeros of $\Delta_{\pm}^{-1}$ in the HTL approximation in quadratic scales.

The appearance of thermal masses presents unavoidable problems at some order of perturbation theory, namely when these higher orders probe the energy/momentum scale $k_0, k \sim gT$, and, in particular in the bosonic sector, the Bose-Einstein distribution factors enhance the sensitivity to the infrared.

It is therefore mandatory to switch from bare perturbation theory to one that uses resummed (dressed) propagators. In simple cases like scalar φ^4 theory, where the only HTL is a constant mass term, this resummation has been studied already long ago [29]. Conceptually (although not practically) equally simple is the example of scalar electrodynamics, where the only HTL's are the self energy diagrams considered above. However, already in spinor QED and to a larger extent in QCD, it turns out that there are HTL vertex functions [36] which have to be treated on a par with the HTL self energies to achieve a systematic resummed perturbation theory [23]. Indeed, if N -point vertex functions give rise to HTL's $\propto T^2$, they are as important as bare vertices when the momentum scale is $\sim gT$:

$$\Gamma_{,N}^{\text{HTL}} \sim g^N T^2 k^{2-N} \sim g^{N-2} k^{4-N} \sim \left. \frac{\partial^N \mathcal{L}_{\text{cl.}}}{\partial A^N} \right|_{k \sim gT}. \quad (25)$$

In fact, even for N so high that there is no comparable tree-level vertex, such vertex functions have to be resummed. Already in spinor QED, there exist one-loop HTL vertex functions with two external fermion lines and an arbitrary number of gauge boson lines. In QCD, there are even more HTL vertex functions without tree-level analogue. It turns out that one-loop diagrams with an arbitrary number of external gauge boson lines are HTL.

4.1 HTL effective actions

The HTL resummation programme can be understood as the transition from the fundamental Lagrangian to an effective one generated by ‘integrating out’ the hard momentum modes $k_0, k \sim T$ in one-loop order.

In scalar φ^4 theory, this effective theory differs from the bare one only in a simple thermal mass term, $\mathcal{L}_{\text{scalar}}^{\text{HTL}} = -\frac{1}{2}\mu^2\varphi^2$.

For gauge bosons and fermions, the effective theory is necessarily a non-local one as gauge invariance forbids a simple local thermal mass term and in the case of fermions it is due to the fact that HTL’s do not spoil chiral symmetries. Remarkably, the infinitely many non-local vertex functions can be summarized by a comparatively simple and manifestly gauge-invariant integral representation [77, 25, 37]

$$\begin{aligned}\mathcal{L}^{\text{HTL}} = & \hat{M}^2 \int \frac{d\Omega_v}{4\pi} \bar{\psi} \gamma^\mu \frac{v_\mu}{v \cdot D(A)} \psi \\ & - \frac{3}{2} \omega_{pl.}^2 \text{tr} \int \frac{d\Omega_v}{4\pi} F^{\mu\alpha} \frac{v_\alpha v^\beta}{(v \cdot D_{adj.}(A))^2} F_{\mu\beta}\end{aligned}\quad (26)$$

$v = (1, \vec{v})$ is a light-like 4-vector, i.e. with $\vec{v}^2 = 1$, and its spatial components are averaged over by $d\Omega_v$. v is the remnant of the hard plasma constituents’ momenta $p^\mu \sim T v^\mu$, namely their light-like 4-velocity, and the overall scale T has combined with the coupling constant to form the scale of thermal masses, $\hat{M}, \omega_{pl.} \sim gT$.

The covariant derivatives in the denominators of (26) are responsible for the fact that there are infinitely many HTL’s involving external fermions and an arbitrary number of gauge bosons, even in QED, where only the pure gauge-field sector becomes bilinear because of $D_{adj.}(A) \rightarrow \partial$.

Technically, resummed perturbation theory amounts to the replacement

$$\mathcal{L}_{\text{cl.}} \rightarrow \mathcal{L}_{\text{cl.}} + \mathcal{L}^{\text{HTL}} - \ell \mathcal{L}^{\text{HTL}} \quad (27)$$

where ℓ is a loop counting parameter that is sent to 1 in the end, after the last term has been treated as a ‘thermal counterterm’.

Because $\mathcal{L}^{\text{HTL}}$ has been derived under the assumption of soft external momenta, this prescription is in fact only to be followed for soft propagators and vertices [23]. Those involving hard momenta (if present) do not require this resummation, and they can be excluded from this resummation by the introduction of some intermediate scale Λ with $gT \ll \Lambda \ll T$. Complete results have to come out independent of Λ , of course.

4.2 Example: NLO terms in scalar electrodynamics

Massless scalar electrodynamics [54] is a particularly simple toy model as its HTL effective action (26) is bilinear in all fields. The scalar HTL self-energy (19) is moreover a simple mass term, and in order to consider the one-loop corrections to the photon polarization tensor, nothing more is needed. Let us

consider two limiting cases of this to illustrate some important points of the HTL resummation programme.

4.2.1 Debye mass

The Debye mass, i.e. the inverse screening length of electrostatic fields, is determined by the zero of $\Delta_\ell(k_0 = 0, k)$ at imaginary k leading to $m_D^2 = \Pi_{00}(k_0 = 0, \vec{k}^2 = -m_D^2)$. Because at leading order $\hat{\Pi}_{00}(k_0 = 0, k)$ turns out to be independent of k (see (18)), the frequently found definition [45] of m_D^2 as $\hat{\Pi}_{00}(k_0 = 0, k \rightarrow 0)$ happens to be correct, but becomes unphysical in general [71]: beyond LO, only the former, self-consistent definition is renormalization-group invariant and (in nonabelian theories) gauge invariant.

The resummation programme sketched above makes the LO result $\hat{m}_D^2 = e^2 T^2/3$ part of the new lowest-order Lagrangian $\mathcal{L}_{\text{cl.}} + \mathcal{L}^{\text{HTL}}$. The NLO correction therefore is given by one-loop diagrams using this Lagrangian. These are scalar loops which now have massive propagators with thermal mass $\mu^2 = \hat{\Xi} = e^2 T^2/4$ from (19), from which the HTL result $\hat{m}_D^2$ has to be subtracted as thermal counter-term. Indeed, without this subtraction, the LO result would be generated a second time, since for large loop momenta the thermal mass of the scalar is negligible. Because of the subtraction, the one-loop integrals are now receiving their leading contributions from soft loop momenta $k \sim eT$:

$$\begin{aligned} \delta\Pi_{00}(0, q) &= e^2 \sum_K \frac{4k_0^2}{(K^2 - \mu^2)[(K - Q)^2 - \mu^2]} - e^2 \sum_K \frac{2}{K^2 - \mu^2} - \hat{m}_D^2 \\ &= \frac{e^2}{\pi^2} \int_0^\infty dk k^2 \left\{ \frac{n(\sqrt{k^2 + \mu^2})}{\sqrt{k^2 + \mu^2}} \left[1 + \frac{k^2 + \mu^2}{kq} \ln \left| \frac{2k + q}{2k - q} \right| \right] - (\mu \rightarrow 0) \right\} \quad (28) \end{aligned}$$

To obtain the leading contribution, we can replace $n(E) \rightarrow T/E$ which gives

$$\delta\Pi_{00}(0, q) = -\frac{e^2 T}{\pi^2} \int_0^\infty dk \left\{ \frac{\mu^2}{k^2 + \mu^2} + 1 - \frac{k}{q} \ln \left| \frac{2k + q}{2k - q} \right| \right\} + O(e^2 q^2 \ln(T)). \quad (29)$$

The NLO correction to the Debye mass is now given by evaluating this at $q = i\hat{m}_D$. Incidentally, the above integral is q -independent, as can be seen from an integration by parts, which finally gives

$$m_D^2 = \hat{m}_D^2 - \frac{e^2 T \mu}{2\pi} = \frac{e^2 T^2}{3} \left(1 - \frac{3}{4\pi} e \right). \quad (30)$$

Notice that the perturbative result at NLO involves a single power of e and so is *non-analytic* in $\alpha = e^2/(4\pi)$.

The above calculation can actually be simplified by noting [5] that the only terms capable of producing odd powers in e are the $n = 0$ terms in the sums over Matsubara frequencies in (28). For $n \neq 0$, one has $-k_0^2 = (2\pi n)^2 T^2 \gg \mu^2$ so that the thermal masses can be expanded out, leading to powers of e^2 only.

Keeping only the $n = 0$ contributions, the first integral in (28) vanishes, and we have

$$\delta\Pi_{00}(0, q) = 2e^2 T \sigma^{2\epsilon} \int \frac{d^{3-2\epsilon} k}{(2\pi)^{3-2\epsilon}} \frac{1}{k^2 + \mu^2} \xrightarrow{\epsilon \rightarrow 0} -\frac{e^2 T \mu}{2\pi} \quad (31)$$

in agreement with (30). Here we have introduced dimensional regularization (with mass scale σ) to render the integral finite as in Ref. [5].

As we shall see presently, this simplified resummation by *dimensional reduction* is only possible in the static case ($q_0 = 0$), and not for dynamical quantities.

4.2.2 Plasma frequency

Propagating modes exist for frequencies $q_0 \geq \omega_{\text{pl.}}$, and this plasma frequency is the same for the transverse and longitudinal modes, because it corresponds to the long-wavelength limit $q \rightarrow 0$, and with $\vec{q} = 0$ there is no way to distinguish the polarizations. In the HTL approximation, $\omega_{\text{pl.}}^2 = \hat{m}_D^2/3 = e^2 T^2/9$.

The NLO correction in the case of scalar electrodynamics can be calculated in full analogy to (28), but now with $q_0 = \omega_{\text{pl.}}$ and $\vec{q} \rightarrow 0$. Because of $\vec{q} = 0$, the angular integrals are now trivial and one finds

$$\delta\Pi_\ell(q_0, 0) = \delta\Pi_t(q_0, 0) = -\frac{e^2 T}{2\pi} \left\{ \mu + \frac{4}{3q_0^2} \left([\mu^2 - q_0^2/4]^{3/2} - \mu^3 \right) \right\}. \quad (32)$$

Evaluated at $q_0 = \omega_{\text{pl.}}$ this gives $\delta\omega_{\text{pl.}}^2/\omega_{\text{pl.}}^2 = -e(8\sqrt{2} - 9)/(2\pi) \approx -0.37e$. Notice that without resumming μ , the result would have been completely misleading: evaluating the unresummed result, i.e. (32) with $\mu = 0$, at $q_0 = \omega_{\text{pl.}}$ would have given a purely imaginary result that one would have wrongly identified with a damping constant.²

Furthermore, the correct result (32) is now only obtained if the nonstatic modes are resummed along with the static ones. If one keeps only the zero modes and ignores that in the imaginary time formalism the external frequency q_0 has to be a multiple of $2\pi iT$ (and so cannot be soft and nonzero), but immediately ‘continues’ to $q_0 = \omega_{\text{pl.}} \sim eT$, one would find

$$\delta\Pi_{\ell,t}(q_0, 0)|_{\substack{0\text{-mode} \\ \text{contr.}}} = -\frac{e^2 T}{6\pi} \left\{ \frac{2}{q_0} \left[\frac{\mu}{q_0} - \sqrt{\frac{\mu^2}{q_0^2} - 1} \right] (q_0^2 - \mu^2) + \mu \right\} \quad (33)$$

which clearly differs from (32). The resulting $\delta\omega_{\text{pl.}}^2$ would in fact be only about a quarter of the true result. So dynamic quantities require the full HTL resummation method; resumming only the zero modes is not sufficient (see also [31]).

²In scalar electrodynamics, unlike QCD, the plasmon damping is a higher order effect because the scalar HTL quasiparticles are heavier than plasmons and moreover do not have Landau damping cuts in the HTL approximation.

4.3 NLO corrections for QCD quasiparticles

The calculation of NLO corrections to the long-wavelength plasmons in QCD was in fact one of the first applications of the HTL resummation programme. In particular, the damping constant of order $g^2 T \sim g \omega_{\text{pl}}$ was the subject of a long controversy (in particular with regard to its gauge-fixing (in)dependence) before it was calculated in Ref. [24] with the gauge independent³ result

$$\gamma \approx 0.264 \sqrt{N} g \omega_{\text{pl}}. \quad (34)$$

(for pure-gluon QCD). The analogous calculation for the damping constant of long-wavelength fermionic quasiparticles was carried out in Refs. [50, 26]. The significance of these results is that gluonic and fermionic quasiparticles, which in the HTL approximation appear to be stable, experience damping. For $g \ll 1$, they are weakly damped, whereas for $g \sim 1$ which is more relevant for experimentally accessible quark-gluon plasmas, the damping is significant: $\gamma \sim \frac{1}{2} \omega_{\text{pl}}$.

The NLO correction to the gluonic plasma frequency has also been calculated [76] with the result $\delta \omega_{\text{pl}}^2 / \omega_{\text{pl}}^2 \approx -0.18 \sqrt{N} g$.

The NLO correction to the Debye mass, however, runs into IR problems. Naively one would expect problems from the masslessness of magnetostatic gluons only at two-loop order resummed perturbation theory, when their self-interactions become relevant. However, because gauge independence requires evaluation on mass-shell (which in the case of the Debye mass means $q_0 = 0$, $\vec{q}^2 = -\hat{m}_D^2$), there appear ‘mass-shell singularities’ caused by the massless magnetostatic modes. Because these singularities are only logarithmic, the leading log is perturbatively calculable and reads [71, 72]

$$\delta m_D^2 / \hat{m}_D^2 = \frac{N}{2\pi} \sqrt{\frac{6}{2N + N_f}} g \log \frac{1}{g} + \mathcal{O}(g). \quad (35)$$

For small coupling, the logarithm dominates over the non-perturbative constant behind the logarithm, and thus the perturbative prediction is that of a *positive* correction to the screening mass. Indeed, lattice simulations of both (gauge-fixed) chromo-electrostatic propagators [28] and gauge-invariant lattice definitions of the nonabelian Debye mass [6, 55] give significant positive corrections to the HTL value.⁴

Such a logarithmic sensitivity to the nonperturbative physics of the chromo-magnetostatic sector has in fact been encountered earlier in the damping of hard

³In fact, later investigations using covariant gauges encountered again gauge dependences [9, 10], which are removed, however, when an infrared cut-off is retained while taking the on-mass-shell limit [70]. This means that there is an infrared singularity in the residue of the pole, which is gauge dependent, but the position of the singularity (no longer a simple pole) is gauge independent in accordance with the theorems of Refs. [51, 52].

⁴These positive corrections are so large in fact that the nonperturbative contributions at order g dominate over the HTL result in the gauge-invariant lattice definitions. When extracted from lattice propagators, which also give gauge-independent results [28], these nonperturbative contributions are considerably smaller and about 1/3 of them can be accounted for by one-loop resummed perturbation theory if one introduces a simple phenomenological magnetic screening mass taken from the lattice [72, 74].

excitations [66, 58, 69]. More generally, it arises for all propagating modes [67] as well as for all finite screening lengths [33]. For nonzero wave-vector $\vec{q}$, only the real corrections to the dispersion law $\omega = \omega(q)$ turn out to be IR safe in one-loop resummed perturbation theory.

5 HTL-resummed thermodynamics

In the previous section we have seen that dynamic quantities cannot be treated by the simplified resummation scheme that resums only static modes. For static quantities like the thermodynamic potential, however, such a resummation works in the sense that it gives a scheme to systematically compute the series expansion in powers (and log's) of the coupling. This calculation has been performed to order $\alpha_s^{5/2}$ in QCD [7] with the result (for pure glue)

$$P = -\Omega/V = \frac{8\pi^2 T^4}{45} \left[1 - \frac{15\alpha_s}{4\pi} + 30\left(\frac{\alpha}{\pi}\right)^{3/2} + \frac{135}{2}\left(\frac{\alpha}{\pi}\right)^2 \log \frac{\alpha}{\pi} \right. \\ \left. - \frac{165}{8} \left(\log \frac{\bar{\mu}}{2\pi T} - 11.49 \right) \left(\frac{\alpha}{\pi}\right)^2 + \frac{495}{2} \left(\log \frac{\bar{\mu}}{2\pi T} - 3.23 \right) \left(\frac{\alpha}{\pi}\right)^{5/2} + \dots \right]. \quad (36)$$

Unfortunately, this is very poorly convergent: only when $\alpha_s < 0.05$ one has apparent convergence, but this corresponds to temperatures higher than $10^5 T_c$!

In what follows we shall attempt a different route that resums also the non-static modes, and tries to keep resummation effects even when they are formally of higher order than that achievable at a given loop order.

5.1 Screened perturbation theory

In scalar φ^4 theory, it has been shown [46, 4] that the convergence of thermal perturbation theory can be improved if the thermal mass of the scalar quasi-particles is kept within thermal integrals and not treated as proportional to a coupling constant when setting up the perturbation series. Technically, this is just as in (27), but without the requirement that the resummation has to take place in soft quantities only. Because this changes the UV structure at any finite order of perturbation theory, this introduces new UV divergences and associated renormalization scheme dependences, which in principle can become arbitrarily large. But starting from two-loop order, these can be minimized if the thermal mass used in screened perturbation theory is determined by a variational principle, i.e. a principle of ‘minimal sensitivity’ to the mass parameter used in this reorganization of perturbation theory [4, 73].

In Refs. [2, 3], this approach has been adapted to a one-loop calculation of the thermodynamic potential of QCD where in place of a simple mass term the gauge-invariant HTL effective action is used. While the leading-order interaction term $\propto g^2$ is incomplete (in fact, it is over-included), it does contain the plasmon term $\propto g^3$ without leading to the disastrous result of a thermodynamic pressure

in excess of the Stefan-Boltzmann limit. However, it remains to be seen if this is still true in the (technically very difficult) two-loop approximation, which presumably has to contribute with positive sign to make up for the over-included negative leading-order interaction term of the one-loop approximation.

In the following, I shall present instead the result of a different strategy. Instead of using the HTL effective action as a gauge invariant mass term for an optimization of perturbation theory, which does not (have to) care whether the HTL effective action remains accurate for hard momenta (which it does not), the formalism of self-consistent “ Φ -derivable” [12] approximations will be invoked to find expressions that keep resummation effects for both soft and hard momenta. As we shall see, the leading-order effects will arise exclusively from kinematical regimes where the HTL approximation remains justifiable. Moreover, it will be possible to avoid the spurious UV problems of screened perturbation theory.

5.2 Approximately self-consistent resummations

In the Luttinger-Ward representation of the thermodynamic potential $\Omega = -PV$ [62] (to particle physicists often more familiar as the composite operator effective potential [27]) is expressed as a functional of full propagators D and two-particle irreducible diagrams. Considering for simplicity a scalar field theory for the moment, $\Omega[D]$ has the form

$$\begin{aligned}\Omega[D] &= -T \log Z = \frac{1}{2}T \text{Tr} \log D^{-1} - \frac{1}{2}T \text{Tr} \Pi D + T\Phi[D] \\ &= \int \frac{d^4k}{(2\pi)^4} n(\omega) \text{Im} [\log D^{-1}(\omega, k) - \Pi(\omega, k)D(\omega, k)] + T\Phi[D],\end{aligned}\quad (37)$$

where Tr denotes the trace in configuration space, and $\Phi[D]$ is the sum of the 2-particle-irreducible “skeleton” diagrams

$$-\Phi[D] = \frac{1}{12} \text{---}\bigcirc\text{---} + \frac{1}{8} \text{---}\bigcirc\text{---}\bigcirc\text{---} + \frac{1}{48} \text{---}\bigcirc\text{---}\bigcirc\text{---}\bigcirc\text{---} + \dots \quad (38)$$

The self energy $\Pi = D^{-1} - D_0^{-1}$, where D_0 is the bare propagator, is related to $\Phi[D]$ by

$$\delta\Phi[D]/\delta D = \frac{1}{2}\Pi. \quad (39)$$

An important property of the functional $\Omega[D]$, which is easily verified using (39), is that it is stationary under variations of D :

$$\delta\Omega[D]/\delta D = 0. \quad (40)$$

Self-consistent (“ Φ -derivable”) [12] approximations are obtained by selecting a class of skeletons in $\Phi[D]$ and calculating Π from Eq. (39) above, preserving the stationarity condition.

The stationarity of $\Omega[D]$ has an interesting consequence for the entropy $\mathcal{S} = -\partial(\Omega/V)/\partial T$. Because of Eq. (40), the temperature derivative of the spectral density in the dressed propagator cancels out and only the explicit

Bose-Einstein factors need to be differentiated in (37), yielding [75, 78, 15, 17]

$$\begin{aligned} \mathcal{S} = & - \int \frac{d^4 k}{(2\pi)^4} \frac{\partial n(\omega)}{\partial T} \text{Im} \log D^{-1}(\omega, k) \\ & + \int \frac{d^4 k}{(2\pi)^4} \frac{\partial n(\omega)}{\partial T} \text{Im} \Pi(\omega, k) \text{Re} D(\omega, k) + \mathcal{S}' \end{aligned} \quad (41)$$

with

$$\mathcal{S}' = - \frac{\partial(T\Phi)}{\partial T} \Big|_D + \int \frac{d^4 k}{(2\pi)^4} \frac{\partial n(\omega)}{\partial T} \text{Re} \Pi \text{Im} D = 0 \quad (42)$$

up to terms that are of loop-order 3 or higher. Thus, in contrast to Ω , where Φ contributes already to order g^2 in perturbation theory, Eq. (5) with $\mathcal{S}' = 0$ is perturbatively correct to order g^3 . The first two terms in Eq. (41) represent essentially the entropy of “independent quasiparticles”, while $\mathcal{S}'$ may be viewed as the residual interactions among these quasiparticles [78].

The same simplification holds true in the presence of fermions and, with nonzero chemical potential, extends to the fermion density $\mathcal{N} = -\partial(\Omega/V)/\partial\mu$ [16, 17]. In a *self-consistent* two-loop approximation one thus has the remarkably simple formulae, now for general theories

$$\begin{aligned} \mathcal{S} = & -\text{tr} \int \frac{d^4 k}{(2\pi)^4} \frac{\partial n(\omega)}{\partial T} [\text{Im} \log D^{-1}(\omega, k) - \text{Im} \Pi(\omega, k) \text{Re} D(\omega, k)] \\ & - 2 \text{tr} \int \frac{d^4 k}{(2\pi)^4} \frac{\partial f(\omega)}{\partial T} [\text{Im} \log S^{-1}(\omega, k) - \text{Im} \Sigma(\omega, k) \text{Re} S(\omega, k)] , \end{aligned} \quad (43)$$

$$\mathcal{N} = - 2 \text{tr} \int \frac{d^4 k}{(2\pi)^4} \frac{\partial f(\omega)}{\partial \mu} [\text{Im} \log S^{-1}(\omega, k) - \text{Im} \Sigma(\omega, k) \text{Re} S(\omega, k)] , \quad (44)$$

where $n(\omega) = (e^{\beta\omega} - 1)^{-1}$, $f(\omega) = (e^{\beta(\omega - \mu)} + 1)^{-1}$, and “tr” refers to all discrete labels, including spin, color and flavor when applicable.

In gauge theories, the above expressions have to be augmented by Faddeev-Popov ghost contributions which enter like bosonic fields but with opposite overall sign, unless a gauge is used where the ghosts do not propagate such as in axial gauges. But because Φ -derivable approximations do not generally respect gauge invariance,⁵ the self-consistent two-loop approximation will not be gauge-fixing independent. It is in fact not even clear that the corresponding gap equations (39) have solutions at all or that one can renormalize these (nonperturbative) equations.

For this reason, we shall construct *approximately self-consistent* solutions which are gauge invariant and which maintain equivalence with conventional perturbation theory up to and including order g^3 , the maximum (perturbative) accuracy of the two-loop approximation for Φ . For these approximations it will be sufficient to keep only the two transverse structure functions of the gluon propagator and to neglect ghosts.

⁵For this, one would have to treat vertices on an equal footing with self-energies, which is in principle possible using the formalism developed in Ref. [35].

For soft momenta, we know that the leading order contribution is given by the HTL's, and indeed there is no HTL ghost self-energy. For hard momenta, one can identify the contributions to (43) below order g^4 as those linear in the self-energies,

$$\begin{aligned}\mathcal{S}^{\text{hard}} &= \mathcal{S}_0 + 2N_g \int \frac{d^4k}{(2\pi)^4} \frac{\partial n}{\partial T} \text{Re} \Pi_t \text{Im} \frac{1}{\omega^2 - k^2} \\ &- 4N_f \int \frac{d^4k}{(2\pi)^4} \frac{\partial f}{\partial T} \left\{ \text{Re} \Sigma_+ \text{Im} \frac{-1}{\omega - k} - \text{Re} \Sigma_- \text{Im} \frac{-1}{\omega + k} \right\}\end{aligned}\quad (45)$$

considering now a gauge theory with N_g gluons and N_f fermion flavors. Because the imaginary parts of the free propagators restrict their contribution to the light-cone, only the light-cone projections of the self-energies enter. At order g^2 this is exactly given by the HTL results, without having to assume soft ω, k [53, 32] (as in the above example of scalar electrodynamics, see Eqs. (15), (19))

$$\text{Re} \Pi_t^{(2)}(\omega^2 = k^2) = \hat{\Pi}_t(\omega^2 = k^2) = \frac{1}{2} \hat{m}_D^2 \equiv m_\infty^2, \quad (46)$$

$$2k \text{Re} \Sigma_\pm^{(2)}(\omega = \pm k) = 2k \hat{\Sigma}_\pm(\omega = \pm k) = 2\hat{M}^2 \equiv M_\infty^2, \quad (47)$$

and without contributions from the other components of $\Pi_{\mu\nu}$ and the Faddeev-Popov self-energy.

There is no contribution $\propto g^2$ from soft momenta in (43) and (44) so that one is left with remarkably simple general formulae for the leading-order interaction contributions to the thermodynamic potentials expressed through the asymptotic thermal masses of the bosonic and fermionic quasiparticles:

$$\mathcal{S}^{(2)} = -T \left\{ \sum_B \frac{m_{\infty B}^2}{12} + \sum_F \frac{M_{\infty F}^2}{24} \right\}, \quad \mathcal{N}^{(2)} = -\frac{1}{8\pi^2} \sum_F \mu_F M_{\infty F}^2. \quad (48)$$

Here the sums run over all the bosonic (B) and fermionic (F) degrees of freedom (e.g. 4 for each Dirac fermion), which are allowed to have different asymptotic masses and, in the case of fermions, different chemical potentials.

Turning now to the next order, g^3 , let us first recapitulate how this usually arises in the thermodynamic potential. Since this is a static quantity, we can use the imaginary-time formalism and concentrate on the zero-modes as in Sect. 4.2.1. For $\omega = 0$ the leading contributions to the self-energies are $\hat{\Pi}_\ell(0, k) = \hat{m}_D^2 \propto (gT)^2$ and $\hat{\Pi}_t(0, k) = 0$. This gives the “plasmon-effect”⁶ term

$$P^{(3)} = -N_g T \int \frac{d^3k}{(2\pi)^3} \left[\log \left(1 + \frac{\hat{m}_D^2}{k^2} \right) - \frac{\hat{m}_D^2}{k^2} \right] = N_g \frac{\hat{m}_D^3 T}{12\pi}. \quad (49)$$

In deriving the self-consistent expressions for entropy and density, Eqs. (43) and (44), we can no longer use this argument to extract the order g^3 term, for we have first rewritten the thermodynamic potential in terms of real-time

⁶Obviously a misnomer. It is caused exclusively by the Debye screening mass, not the plasmon mass = plasma frequency $\neq \hat{m}_D$.

propagators and self-energies and then used stationarity to drop T and μ derivatives of the spectral densities hidden in the full propagators. In fact, from the second line of (37), one can still isolate the zero-mode contribution (49) by $\int \frac{d\omega}{\pi\omega} \text{Im}[\dots] = [\dots](\omega = 0)$, but in (43) and (44) we have products of imaginary and real parts times statistical distribution functions.

Indeed, the order g^3 contributions now arise from both soft and hard momentum regimes. In $\mathcal{S}^{\text{soft}}$ such contributions are due to the singular behaviour of $\partial n/\partial T \sim 1/\omega$ which does not allow us to expand out the self-energy insertions perturbatively, but on dimensional grounds gives a contribution $\sim \Pi^{3/2}$. In $\mathcal{S}^{\text{hard}}$, on the other hand, where we could expand out the self-energies as in Eq. (45), g^3 contributions arise from NLO order contributions to Π and Σ themselves. (In the case of the pressure, this did not happen because of the stationarity property of the pressure [17].)

With this insight, we can now formulate an *approximately self-consistent* dressing of the propagators that is in line with the maximum perturbative accuracy of the Φ -derivable two-loop approximation: for soft momenta, we take the (gauge-invariant and gauge-independent) HTL expressions for self-energies and propagators; for hard momenta, where according to (45) only the light-cone limit of the self-energies contribute below order g^4 , the correct leading-order contribution is still given by the HTL expressions (46), (47), but in order to include the g^3 contributions completely, we also require the NLO corrections to the on-light-cone self-energies. The latter can be calculated by standard HTL perturbation theory, and the theorems of Ref. [52] ensure their gauge independence.

5.2.1 HTL approximation

As a first approximation let us consider one which only uses the HTL expressions without NLO corrections thereof. We have seen that this gives the correct leading-order interaction term $\propto g^2$, some part of the g^3 contribution, and infinitely many formally higher-order terms as well, since we are going to use (43) and (44) “non-perturbatively”, i.e. without expanding out in powers of g and truncating. We can do so because expressions (43) and (44) are manifestly UV finite, for they involve only the derivatives of the statistical distribution functions—in (37), the $T = 0$ UV-divergences are contained in the integration domain $\omega \rightarrow -\infty$, where the undifferentiated n ’s and f ’s do not fall off exponentially.

Using the HTL expressions in (43) and considering for simplicity the pure-gluon case, we obtain two physically distinct contributions. The first corresponds to the transverse and longitudinal gluonic quasiparticle poles,

$$\mathcal{S}_{\text{HTL}}^{\text{QP}} = -N_g \int \frac{k^2 dk}{2\pi^2} \frac{\partial}{\partial T} \left[2T \log(1 - e^{-\omega_t(k)/T}) + T \log \frac{1 - e^{-\omega_\ell(k)/T}}{1 - e^{-k/T}} \right], \quad (50)$$

where only the explicit T dependences are to be differentiated, and not those implicit in the HTL dispersion laws $\omega_t(k)$ and $\omega_\ell(k)$. Secondly, there are the

Landau-damping contributions which read

$$\begin{aligned} \mathcal{S}_{\text{HTL}}^{\text{LD}} = & -N_g \int_0^\infty \frac{k^2 dk}{2\pi^3} \int_0^k d\omega \frac{\partial n(\omega)}{\partial T} \left\{ 2 \arg[k^2 - \omega^2 + \hat{\Pi}_t] \right. \\ & \left. - 2 \text{Im} \hat{\Pi}_t \text{Re} [\omega^2 - k^2 - \hat{\Pi}_t]^{-1} + \arg[k^2 + \hat{\Pi}_\ell] - \text{Im} \hat{\Pi}_\ell \text{Re} [k^2 + \hat{\Pi}_\ell]^{-1} \right\}. \end{aligned} \quad (51)$$

The usual perturbative g^2 -contribution (48) is contained in the first term of Eq. (50); all the other terms in Eqs. (50),(51) are of order g^3 in a small- g expansion.

In the HTL approximation, only the soft plasmon effect $\sim g^3$ contained in $\mathcal{S}^{\text{soft}}$ is present, which turns out to equal

$$\mathcal{S}_3^{\text{soft}} = \frac{\partial P^{(3)}}{\partial T} \Big|_{\hat{m}_D}, \quad (52)$$

which is $\frac{1}{4}\mathcal{S}_3$ in the case of pure glue. However, this identification requires a peculiar sum rule

$$\begin{aligned} \Delta \mathcal{S}_3 & \equiv N_g \int \frac{d^4 k}{(2\pi)^4} \frac{1}{\omega} \left\{ 2 \text{Im} \hat{\Pi}_t \text{Re} (\hat{D}_t - D_t^{(0)}) - \text{Im} \hat{\Pi}_\ell \text{Re} (\hat{D}_\ell - D_\ell^{(0)}) \right\} \\ & \equiv \Delta \mathcal{S}_t^{(3)} + \Delta \mathcal{S}_\ell^{(3)} = 0, \end{aligned} \quad (53)$$

which we found to hold numerically by cancellations in more than 8 significant digits. Rather unusually (cp. Ref. [57]), this does not hold separately for the longitudinal and transverse sector and moreover holds only after carrying out both, the frequency and the momentum integrations in (53).

Although the g^3 term in $\mathcal{S}_{\text{HTL}}$ is only a fraction of the full one, it would make similar troubles when expanded out perturbatively, throwing away all higher-order terms in g : for large enough g , the perturbative approximation would lead to an entropy in excess of the Stefan-Boltzmann limit. $\mathcal{S}_{\text{HTL}}$, on the other hand, is a monotonically decreasing function of $\hat{m}_D/T \sim g$.

5.2.2 Next-to-leading approximation

The plasmon term $\sim g^3$ becomes complete only upon inclusion of the next-to-leading correction to the asymptotic thermal masses m_∞ and M_∞ . These are determined in standard HTL perturbation theory through

$$\begin{aligned} \delta m_\infty^2(k) &= \text{Re} \delta \Pi_T(\omega = k) \\ &= \text{Re} (\text{blob on wavy line} + \text{blob on thick dashed line} + \text{blob on wavy line} + \text{blob on thick dashed line})|_{\omega=k} \end{aligned} \quad (54)$$

where thick dashed and wiggly lines with a blob represent HTL propagators for longitudinal and transverse polarizations, respectively. Similarly,

$$\frac{1}{2k} \delta M_\infty^2(k) = \delta \Sigma_+(\omega = k) = \text{Re} (\text{blob on thick dashed line} + \text{blob on wavy line})|_{\omega=k}. \quad (55)$$

The explicit proof that these contributions indeed restore the correct plasmon term is given in Ref. [17].

These corrections to the asymptotic thermal masses are, in contrast to the latter, nontrivial functions of the momentum, which can be evaluated only numerically. However, as far as the generation of the plasmon term is concerned, these functions contribute in the averaged form

$$\bar{\delta}m_\infty^2 = \frac{\int dk k n'_{\text{BE}}(k) \text{Re } \delta\Pi_T(\omega = k)}{\int dk k n'_{\text{BE}}(k)} \quad (56)$$

(cp. Eq. (48)) and similarly

$$\bar{\delta}M_\infty^2 = \frac{\int dk k n'_{\text{FD}}(k) \text{Re } 2k\delta\Sigma_+(\omega = k)}{\int dk k n'_{\text{FD}}(k)}. \quad (57)$$

These averaged asymptotic thermal masses turn out to be given by the remarkably simple expressions [17]

$$\bar{\delta}m_\infty^2 = -\frac{1}{2\pi}g^2NT\hat{m}_D, \quad \bar{\delta}M_\infty^2 = -\frac{1}{2\pi}g^2C_fT\hat{m}_D, \quad (58)$$

where $C_f = N_g/(2N)$. Since the integrals in (56) and (57) are dominated by hard momenta, these thermal mass corrections only pertain to hard excitations. Indeed, in Sect. 4.3 we have seen that e.g. the plasmon mass at $k = 0$ receives a different, namely smaller correction, whereas the NLO contribution to the Debye mass is even positive and logarithmically enhanced.

Pending the full evaluation of the NLO corrections to $\text{Re } \delta\Pi$ and $\text{Re } \delta\Sigma$, it has therefore been proposed in Refs. [15, 16, 17] to define a next-to-leading approximation through (for gluons)

$$\mathcal{S}_{NLA} = \mathcal{S}_{HTL}\Big|_{\text{soft}} + \mathcal{S}_{\bar{m}_\infty^2}\Big|_{\text{hard}}, \quad (59)$$

where $\bar{m}_\infty^2$ includes (58). To separate soft ($k \sim \hat{m}_D$) and hard ($k \sim 2\pi T$) momentum scales, we introduce the intermediate scale $\Lambda = \sqrt{2\pi T\hat{m}_D c_\Lambda}$ and consider a variation of $c_\Lambda = \frac{1}{2} \dots 2$ as part of our theoretical uncertainty.

Another crucial issue concerns the definition of the corrected asymptotic mass $\bar{m}_\infty$. For the range of coupling constants of interest ($g \gtrsim 1$), the correction $|\bar{\delta}m_\infty^2|$ is greater than the LO value m_∞^2 , leading to tachyonic masses if included in a strictly perturbative manner.

However, this problem is not at all specific to QCD. In the simple $g^2\varphi^4$ model, one-loop resummed perturbation theory gives

$$m^2 = g^2T^2\left(1 - \frac{3}{\pi}g\right) \quad (60)$$

which also turns tachyonic for $g \gtrsim 1$. On the other hand, the self-consistent solution of the corresponding one-loop gap equation [29, 30]

$$m^2 = \Pi(m) = 12g^2 \sum_K \frac{-1}{K^2 - m^2} \quad (61)$$




Figure 3: Comparison of the lattice data for the entropy of pure-gluon SU(3) gauge theory of Ref. [19] (gray band) with the range of $\mathcal{S}_{HTL}$ (solid lines) and $\mathcal{S}_{NLA}$ (dash-dotted lines) for $\bar{\mu} = \pi T \dots 4\pi T$ and $c_\Lambda = 1/2 \dots 2$.

(properly renormalized), whose perturbative expansion begins exactly like (60), is a monotonic function in g . In fact, it turns out that the first two terms in a (m/T) -expansion of this gap equation,

$$m^2 = g^2 T^2 - \frac{3}{\pi} g^2 T m, \quad (62)$$

which is perturbatively equivalent to (60), has a solution that is extremely close to that of the full gap equation (for $\overline{\text{MS}}$ renormalization scales $\bar{\mu} \approx 2\pi T$) [17].

In QCD, the (non-local) gap equations are way too complicated to be attacked directly. We instead consider perturbatively equivalent expressions for the corrected $\bar{m}_\infty$ which are monotonic functions in g . Besides the solution to a quadratic equation analogous to (62) we have tried the simplest Padé approximant $m^2 = g^2 T^2 / (1 + \frac{3}{\pi} g)$, which also gives a greatly improved approximation to the solution of scalar gap equations. In QCD, our final results do not depend too much on whether we use the Padé approximant [15, 16] or a quadratic gap equation [17].

The main uncertainty rather comes from the choice of the renormalization scale which determines the magnitude of the strong coupling constant when this is taken as determined by the renormalization group equation (2-loop in the following).

In Fig. 3, the numerical results for the HTL entropy and the NLA one are given as a function of T/T_c with T_c chosen as $T_c = 1.14\Lambda_{\overline{\text{MS}}}$. The full lines show the range of results for $\mathcal{S}_{HTL}$ when the renormalization scale $\bar{\mu}$ is varied from πT to $4\pi T$; the dash-dotted lines mark the corresponding results for $\mathcal{S}_{NLA}$ with the additional variation of c_Λ from $1/2$ to 2 . The dark-gray band are lattice data from Ref. [19]. Evidently, there is very good agreement for $T \gtrsim 2.5T_c$.

From the above results for the entropy density, one can recover the thermodynamic pressure by simple integration, $P(T) - P(T_0) = \int_{T_0}^T dT' S(T')$. The integration constant $P(T_0)$, however, is a strictly nonperturbative input. It cannot be fixed by requiring $P(T=0) = 0$, as this is in the confinement regime. It is also not sufficient to know that $\lim_{T \rightarrow \infty} P = P_{\text{free}}$ by asymptotic freedom. In fact, the undetermined integration constant in $P(T)/P_{\text{free}}(T)$ when expressed as a function of $\alpha_s(T)$ corresponds to a term [16] $C \exp\{-\alpha_s^{-1}[4\beta_0^{-1} + O(\alpha_s)]\}$, which vanishes for $\alpha_s \rightarrow 0$ with all derivatives and thus is not fixed by any order of perturbation theory. It is, in essence, the nonperturbative bag constant, which can be added on to standard perturbative results, too. However, in $P(T)/P_{\text{free}}(T)$ this term becomes rapidly unimportant as the temperature is increased, as it decays like T^{-4} . Fixing it by $P(T_c) = 0$, which is a good approximation in particular for the pure-gluon case because glue balls are rather heavy, one finds again good agreement with lattice data for $T \gtrsim 2.5T_c$.

This approach can be generalized [16, 17] also to nonzero chemical potentials μ_f , where lattice data are not available⁷. Simpler quasiparticle models [64, 59] have already been used to extrapolate lattice data to finite chemical potential [65]. The HTL approach offers a possible refinement, but that has still to be worked out.

6 Conclusion

Hard thermal loops, the leading-order contributions to self-energies and vertices at high temperature and/or density, form a gauge-invariant basis for a systematic perturbative expansion, as long as one does not run into the perturbative barrier formed by the inherently nonperturbative sector of self-interacting chromomagnetostatic modes. But in QCD one faces the additional problem that corrections to leading-order results are so large for almost all values of the coupling of interest that they lead to a complete loss of (apparent) convergence. However, we have seen that further resummations which keep as much as possible of the effects of HTL resummation without expanding in a power series in the coupling may lead to results that remain valid down to temperatures a few times the deconfinement phase transition temperature.

⁷Lattice results exist however for the response of thermodynamic quantities to infinitesimal chemical potentials, namely for quark number susceptibilities [41, 38, 39], and the above approach has been applied to those, too, by now [18].

References

- [1] Akhiezer, I. A. and S. V. Peletminskii: 1960. *Sov. Phys. JETP* **11**, 1316.
- [2] Andersen, J. O., E. Braaten, and M. Strickland: 1999. *Phys. Rev. Lett.* **83**, 2139.
- [3] Andersen, J. O., E. Braaten, and M. Strickland: 2000. *Phys. Rev.* **D61**, 014017, 074016.
- [4] Andersen, J. O., E. Braaten, and M. Strickland: 2001. *Phys. Rev.* **D63**, 105008.
- [5] Arnold, P. and O. Espinosa: 1993. *Phys. Rev.* **D47**, 3546.
- [6] Arnold, P. and L. G. Yaffe: 1995. *Phys. Rev.* **D52**, 7208.
- [7] Arnold, P. and C.-X. Zhai: 1995. *Phys. Rev.* **D51**, 1906.
- [8] Aurenche, P., F. Gelis, R. Kobes, and E. Petitgirard: 1997. *Z. Phys.* **C75**, 315.
- [9] Baier, R., G. Kunstatter, and D. Schiff: 1992a. *Phys. Rev.* **D45**, 4381.
- [10] Baier, R., G. Kunstatter, and D. Schiff: 1992b. *Nucl. Phys.* **B388**, 287.
- [11] Baier, R., S. Peigné, and D. Schiff: 1994. *Z. Phys.* **C62**, 337.
- [12] Baym, G.: 1962. *Phys. Rev.* **127**, 1391.
- [13] Blaizot, J.-P. and E. Iancu: 2001, ‘The quark-gluon plasma: Collective dynamics and hard thermal loops’, hep-ph/0101103.
- [14] Blaizot, J.-P., E. Iancu, and R. R. Parwani: 1995. *Phys. Rev.* **D52**, 2543.
- [15] Blaizot, J.-P., E. Iancu, and A. Rebhan: 1999a. *Phys. Rev. Lett.* **83**, 2906.
- [16] Blaizot, J.-P., E. Iancu, and A. Rebhan: 1999b. *Phys. Lett.* **B470**, 181.
- [17] Blaizot, J.-P., E. Iancu, and A. Rebhan: 2001a. *Phys. Rev.* **D63**, 065003.
- [18] Blaizot, J.-P., E. Iancu, and A. Rebhan: 2001b, ‘Quark number susceptibilities from HTL-resummed thermodynamics’, hep-ph/0110369.
- [19] Boyd, G., J. Engels, F. Karsch, E. Laermann, C. Legeland, M. Lütgemeier, and B. Petersson: 1996. *Nucl. Phys.* **B469**, 419.
- [20] Braaten, E.: 1995. *Phys. Rev. Lett.* **74**, 2164.
- [21] Braaten, E. and A. Nieto: 1996a. *Phys. Rev.* **D53**, 3421.
- [22] Braaten, E. and A. Nieto: 1996b. *Phys. Rev. Lett.* **76**, 1417.
- [23] Braaten, E. and R. D. Pisarski: 1990a. *Nucl. Phys.* **B337**, 569.

- [24] Braaten, E. and R. D. Pisarski: 1990b. *Phys. Rev.* **D42**, 2156.
- [25] Braaten, E. and R. D. Pisarski: 1992a. *Phys. Rev.* **D45**, 1827.
- [26] Braaten, E. and R. D. Pisarski: 1992b. *Phys. Rev.* **D46**, 1829.
- [27] Cornwall, J. M., R. Jackiw, and E. Tomboulis: 1974. *Phys. Rev.* **D10**, 2428.
- [28] Cucchieri, A., F. Karsch, and P. Petreczky: 2001. *Phys. Rev.* **D64**, 036001.
- [29] Dolan, L. and R. Jackiw: 1974. *Phys. Rev.* **D9**, 3320.
- [30] Drummond, I. T., R. R. Horgan, P. V. Landshoff, and A. Rebhan: 1998. *Nucl. Phys.* **B524**, 579.
- [31] Elze, H. T., U. Heinz, K. Kajantie, and T. Toimela: 1988. *Z. Phys.* **C37**, 305.
- [32] Flechsig, F. and A. K. Rebhan: 1996. *Nucl. Phys.* **B464**, 279.
- [33] Flechsig, F., A. K. Rebhan, and H. Schulz: 1995. *Phys. Rev.* **D52**, 2994.
- [34] Fradkin, E. S.: 1965. *Proc. Lebedev Inst.* **29**, 1.
- [35] Freedman, B. A. and L. D. McLerran: 1977. *Phys. Rev.* **D16**, 1130.
- [36] Frenkel, J. and J. C. Taylor: 1990. *Nucl. Phys.* **B334**, 199.
- [37] Frenkel, J. and J. C. Taylor: 1992. *Nucl. Phys.* **B374**, 156.
- [38] Gai, R. V. and S. Gupta: 2001. *Phys. Rev.* **D64**, 074506.
- [39] Gai, R. V., S. Gupta, and P. Majumdar: 2001, ‘Susceptibilities and screening masses in two flavor QCD’, hep-lat/0110032.
- [40] Gell-Mann, M. and K. A. Brueckner: 1957. *Phys. Rev.* **106**, 364.
- [41] Gottlieb, S., W. Liu, D. Toussaint, R. L. Renken, and R. L. Sugar: 1987. *Phys. Rev. Lett.* **59**, 2247.
- [42] Gross, D. J., R. D. Pisarski, and L. G. Yaffe: 1981. *Rev. Mod. Phys.* **53**, 43.
- [43] Hatsuda, T.: 1997. *Phys. Rev.* **D56**, 8111.
- [44] Kalashnikov, O. K. and V. V. Klimov: 1980. *Sov. J. Nucl. Phys.* **31**, 699.
- [45] Kapusta, J. I.: 1989, *Finite-temperature field theory*. Cambridge, UK: Cambridge University Press.
- [46] Karsch, F., A. Patkós, and P. Petreczky: 1997. *Phys. Lett.* **B401**, 69.
- [47] Kastening, B.: 1997. *Phys. Rev.* **D56**, 8107.

- [48] Klimov, V. V.: 1981. *Sov. J. Nucl. Phys.* **33**, 934.
- [49] Kobes, R.: 1990. *Phys. Rev.* **D42**, 562.
- [50] Kobes, R., G. Kunstatter, and K. Mak: 1992. *Phys. Rev.* **D45**, 4632.
- [51] Kobes, R., G. Kunstatter, and A. Rebhan: 1990. *Phys. Rev. Lett.* **64**, 2992.
- [52] Kobes, R., G. Kunstatter, and A. Rebhan: 1991. *Nucl. Phys.* **B355**, 1.
- [53] Kraemmer, U., M. Kreuzer, and A. Rebhan: 1990. *Ann. Phys.* **201**, 223 [Appendix].
- [54] Kraemmer, U., A. K. Rebhan, and H. Schulz: 1995. *Ann. Phys.* **238**, 286.
- [55] Laine, M. and O. Philipsen: 1999. *Phys. Lett.* **B459**, 259.
- [56] Landsman, N. P. and C. G. van Weert: 1987. *Phys. Rept.* **145**, 141.
- [57] Le Bellac, M.: 1996, *Thermal Field Theory*. Cambridge, UK: Cambridge University Press.
- [58] Lebedev, V. V. and A. V. Smilga: 1991. *Phys. Lett.* **B253**, 231.
- [59] Levai, P. and U. Heinz: 1998. *Phys. Rev.* **C57**, 1879.
- [60] Lifshitz, E. M. and L. P. Pitaevsky: 1981, *Physical Kinetics*. Oxford, UK: Pergamon Press.
- [61] Linde, A. D.: 1980. *Phys. Lett.* **B96**, 289.
- [62] Luttinger, J. M. and J. C. Ward: 1960. *Phys. Rev.* **118**, 1417.
- [63] Nachbagauer, H., A. K. Rebhan, and D. J. Schwarz: 1996. *Phys. Rev.* **D53**, 882.
- [64] Peshier, A., B. Kämpfer, O. P. Pavlenko, and G. Soff: 1996. *Phys. Rev.* **D54**, 2399.
- [65] Peshier, A., B. Kämpfer, and G. Soff: 2000. *Phys. Rev.* **C61**, 045203.
- [66] Pisarski, R. D.: 1989. *Phys. Rev. Lett.* **63**, 1129.
- [67] Pisarski, R. D.: 1993. *Phys. Rev.* **D47**, 5589.
- [68] Polyakov, A. M.: 1978. *Phys. Lett.* **B72**, 477.
- [69] Rebhan, A.: 1992a. *Phys. Rev.* **D46**, 482.
- [70] Rebhan, A.: 1992b. *Phys. Rev.* **D46**, 4779.
- [71] Rebhan, A. K.: 1993. *Phys. Rev.* **D48**, 3967.
- [72] Rebhan, A. K.: 1994. *Nucl. Phys.* **B430**, 319.

- [73] Rebhan, A.: 2000, ‘Improved resummations for the thermodynamics of the quark gluon plasma’. In: C. P. Korthals Altes (ed.): *Strong and Electroweak Matter 2000*. Singapore: World Scientific Publ., p. 199, hep-ph/0010252.
- [74] Rebhan, A.: 2001, ‘Thermal gauge field theories’, hep-ph/0105183.
- [75] Riedel, E.: 1968. *Z. Phys.* **210**, 403.
- [76] Schulz, H.: 1994. *Nucl. Phys.* **B413**, 353.
- [77] Taylor, J. C. and S. M. H. Wong: 1990. *Nucl. Phys.* **B346**, 115.
- [78] Vanderheyden, B. and G. Baym: 1998. *J. Stat. Phys.* **93**, 843.
- [79] Weldon, H. A.: 1982. *Phys. Rev.* **D26**, 1394.
- [80] Weldon, H. A.: 1989. *Phys. Rev.* **D40**, 2410.